

Study of cosmogenic radiocarbon concentration variations in the Earth's atmosphere during solar activity of Maunder minimum (1645-1715)

S.Tsereteli, V.Bochorishvili, M.Samkharadze, G. Khoriauli

Tbilisi State University, Georgia, 0128. Tbilisi, Chavchavadze av. 3
Email: bochorishvili@icts.tsu.edu.ge

Abstract

Variations of radiocarbon concentration have been studied in annual rings for the last 350 years (1600 – 1950) on the basis of our experimental research using methods of spectral analysis. From this interval of time special attention is paid to the so-called period of the Maunder minimum (1645 – 1715) of solar activity. In the experimental series of corresponding period two types of periodicity are revealed: ≈ 20 year and ≈ 8 year.

1 Introduction

The investigation of the dynamics of solar activity and regularity of its cycling is of great importance for the astrophysics. Correspondingly the study of those intervals when solar activity is sharply changed is of particular significance. Examples of such interesting periods are: 1100-1250 yy "maximum of Middle Ages", in 1282-1342 yy "Volf minimum", 1416-1534 yy "Shperer minimum", 1645-1715 yy "Maunder minimum".

The period of the Maunder minimum has attracted an additional attention because it puts forward a lot of new and unexpected problems. Indeed, for a long time scientists consider spots on the Sun surface as main characteristics of the solar activity. Therefore, the notes about the disappearance of the sunspot is reliable, but due to the absence of regular observations it is difficult to determine level of solar activity in the mentioned period.

It should be noted that the anomalies are found in differential rotation of the Sun surface at the beginning of the Maunder minimum. According of the different authors, it might indicate that global characteristics of the inner parts of the Sun change to such at extant that even the "dynamo" mechanisms is stopped. In this connection the following questions arise: how to characterize the Sun activity and the radioactive state in the Earth's space in that epoch? What is main processes which cause such a drastic fall of solar activity? To what extent our insight into the solar activity correct in general?

For solving mentioned problems physicist direct their special attention to cosmogenic isotopes ^{14}C ($T_{1/2} = 5730 \text{ year}$), ^{10}Be ($T_{1/2} = 1,5 \cdot 10^6 \text{ year}$). The study of these isotopes may give a lot of information about the dynamics of solar activity.

At present the most complete data about the solar activity and the intensity of cosmic rays are obtained by means of the above mentioned cosmogenic isotopes (^{14}C and ^{10}Be) formed in the Earth's atmosphere. The main peculiarity of these isotopes and advantage as compared with others is the fact that there exist objects in nature (e. g. tree rings, Arctic and Antarctic ices, algae with long life time, etc.) which can "fix" i.e. "memorize" annual concentration of the mentioned isotopes, during period that prevails it's half-life time.

2 Radiocarbon Concentration during the Maunder Minimum

To study the influence of solar activity (modulation) in the Earth atmosphere on radiocarbon concentration during the Maunder minimum we use experimental data obtained at the laboratory of nuclear

physics of the TSU [1, 2, 3] (Fig. 1). First of all, it should be noted that experimental data cover the Maunder minimum itself (1645-1720 yy) (Fig. 2) and its neighboring periods (1600-1645 and 1715-1800 yy) too. The measurements in [1] were performed with one-year step, i.e. for every calendar year ^{14}C concentration is measured in the corresponding tree ring. The experimental samples of tree rings were obtained in Lithuania.

The value of ^{14}C concentration in relation to the International standard scale is calculated by the following formula

$$\Delta^{14}\text{C}_0 = \frac{N_s - N_{st}}{N_{st} - N_b},$$

where N_s, N_{st}, N_b represents measured activities of the sample (s), standard (st) and background (b), respectively. Experimental error does not exceed 0.2-0.3%. In order to extract variations of ^{14}C concentration originated only from the "cosmic" sources, it is necessary to exclude the influence of the Earth's magnetic field. This influence has been widely discussed and to date the most accurate estimation is that: global change of geomagnetic field with period of 10 thousand years causes about 10% change in ^{14}C data. The following expression accounts for the geomagnetic field effect in ^{14}C concentration

$$\Delta^{14}\text{C}(t) = \Delta^{14}\text{C}_0(t) - 4,27 - 4,97 \cdot \sin\left[\frac{2\pi}{t + 7388}\right],$$

where t is current calendar year (t is negative B.C.).

As we mentioned above the effective tool for studying the solar activity dynamics during the Maunder minimum is to reveal the cyclings in ^{14}C time dependence (i.e. periodicities in experimental data given in Fig.2). For this purpose the method of spectral analysis has been used [4]. The calculated power spectrum is shown in Fig.3. According to Fig.3 the main peak at frequency 0.05 lies in high reliability interval (95%) (corresponding period ≈ 20 year). Small peak is observed at frequency 0.12 (corresponding period ≈ 8 year) though it's reliability is comparatively low.

3 conclusions:

1. Using the method of spectral analysis we have revealed variations in cosmogenic radiocarbon concentration in the Earth atmosphere during the Maunder minimum.
2. Periodicities of two types have been revealed in experimental data.
3. The reason of the mentioned periodicities is the solar activity. This means that in the Maunder minimum period modulating action of the Sun was weak, nevertheless cycling is preserved. It can be also said that in period of the Maunder minimum the mechanism of solar dynamo still works. Reduction of the number of the sunspots may be caused by the coincidence of the minima of any other long-term periodicities at the moment of the Maunder minimum.

References

- [1] G.E. Kocharov, I.V. Zhorzholiani, Z.V. Lomtadze, R.Y. Metskvarishvili, S.L. Tsereteli, Radiocarbon, 1992, V.34, N2,p. 213-217.
- [2] G.E. Kocharov, Y. N. Markov, R.Y. Metskvarishvili, S. L. Tsereteli. Materials of the Seminar "Intensity of Cosmic Rays and Cosmogen Isotopes". 1983. L. 101.[in Russian]
- [3] I.V. Zhorzholiani, P. G. Kereselidze, et al. Materials of Intern. Seminar "Experimental Methods of Astrophysical and Geophysical Phenomena Study". 1988. L. 114.[in Russian]
- [4] G. M. Jenkins, D. G. Watts " Spectral Analysis and its applications", Holden-Day, San Francisco, Cambridge, London, Amsterdam 1969.
- [5] Catalog of Indices of Solar and Geliomagnetic Activity. Obninsk 1976.[in Russian]




Figure 1: ^{14}C concentration in the atmosphere during 1600-1940 year.




Figure 2: ^{14}C Concentration in the atmosphere during 1600-1800 year.




Figure 3: Dependence of spectral power $\lg(C_{xx})$ on frequency. For radiocarbon concentration in 1645-1715